



# SurFACE™ Sampling Solutions in Food Production



Easy-to-use, sterile specimen collection and surface sampling kits for quality control and identification of hazards at critical control points in food production.

The environment is a principal source of microbiological contamination within a food processing facility. Environmental testing is therefore a crucial part of any food safety program. With a strong focus on environmental testing, Romer Labs provides a broad range of sampling solutions covering **food production**.

## Environmental and Surface Sampling for Food Production Applications

Dry and pre-moistened sterile cellulose sponges and fabric swab are convenient, ready-to-use products. They eliminate the costly and time-consuming steps of diluent preparation, sterilization and application to the sampling devices. They are designed to make it more convenient to sample microbes from surfaces in food processing facilities and industrial settings. The use of different pre-moistening solutions enables a broad range of environmental and product (carcass) sampling applications.

### Available pre-moistening solutions:

**Dey-Engley Neutralizing Broth (DE)** – DE neutralizing broth assists in the recovery of microorganisms in samples taken from surfaces exposed to sanitizing agents. It neutralizes the bactericidal and bacteriostatic effects of sanitizing agents such as chlorine, quaternary ammonium compounds, iodine, phenolics, mercurials, formaldehyde and glutaraldehyde. The medium also contains nutritive agents that aid in recovering and promoting the growth of microorganisms. Bromocresol purple is present in the broth as a colorimetric indicator.

**Neutralizing Buffer (NB)** – The neutralizing buffer assists in the recovery of microorganisms in samples taken from surfaces exposed to sanitizing agents. Of a different color from DE neutralizing broth, it offers the same neutralizing effect but does not include nutrition for the growth of microorganisms. It is therefore classified as a buffer instead of a broth.

**Buffered Peptone Water (BPW)** – Buffered Peptone Water is a pre-enrichment medium useful in isolating *Salmonella* from foods. During many food processes, *Salmonella* and other species can suffer sub-lethal injury from processes involving heat, desiccation, preservatives, pH changes or osmotic pressure shifts. BPW has been shown to facilitate resuscitation of stressed cells and is recommended in the ISO 18593 (Microbiology of the food chain — Horizontal methods for surface sampling) as a moistening solution sampling devices for the recovery of microorganisms from surfaces.

**Lethen Broth (LB)** – Lethen broth is a highly nutritional medium that contains lecithin and Tween 80. It is used for testing the efficacy of sanitizing protocols as lecithin neutralizes quaternary ammonium compounds while Tween 80 neutralizes phenols, hexachlorophene and formalin.

**Maximum Recovery Diluent (MRD)** - Maximum Recovery Diluent is formulated as recommended by ISO 18593 for use as a diluent for general purpose. It is also known as Peptone salt solution (0.1% Peptone, 0.85% NaCl).

## Romer Labs SurfACE™ Sampling Buffers

Ingredient [g/L]	MRD	NB	LB	DE	BPW
Bromocresol purple				0.04	
Casein digest			10	5	
Dextrose				10	
Beef Extract			5		
Lecithin		7	0.7	7	
Neutral Peptone	1				10
Polysorbate 80 (Tween 80)		5	5	5	
Potassium phosphate dibasic		8.5		8	1.5
Sodium Bisulfite		1.25		2.5	
Sodium Bicarbonate					
Sodium Chloride	8.5		5		5
Sodium Phosphate di-basic		3		4	3.5
Sodium Thioglycolate		1		1	
Sodium Thiosulfate		6		6	
Yeast Extract				2.5	

MRD: Maximum Recovery Diluent

NB: Neutralizing Buffer

LB: Letheen Broth

DE: Dey-Engley Neutralizing Broth

BPW: Buffered Peptone Water

## Sponge-Stick

- Sponge-Stick in individual 14.5x30cm (5.5 x 12") Twirl-Tie™ bag
- 7.5 x 4 cm (3 x 1.5") biocide-free cellulose sponge
- Sponge dry or pre-moistened
- Perfectly suited to sample gaps, drains and pipes
- Bags easily opened with attached pull tabs
- Label area for sample identification
- Fast, easy-to-use sponge release mechanism
- Thumb stop helps avoid accidental sponge contact
- Long stick (22 cm, 8.5") for hard-to-reach surfaces
- Easy-to-spot petrol-blue-colored stick
- Less plastic used in production
- No potentially toxic glues used to affix sponge to stick
- Hard plastics with potential to damage bag during transport eliminated
- Compatible with existing extendable poles



### Procedure:



1 Label the bag in the white area. Ballpoint pens can be used. This sampling procedure does not require gloves!




2 Tear bag open by removing the tab. As an added safety measure, the tab is green-striped, making it easy to spot if lost.




3 Pull the bag down from the outside to release the stick. Do not touch the inner surface of the bag.



4 Collect the sample by vigorously rubbing sponge on target area (e.g. 30 x 30 cm, 1 x 1 ft). Do not touch the stick below the thumb-stop located in the middle of the handle.



5 Return the stick to the bag, taking care to insert it no deeper than the thumb-stop.



6 Pull the sponge off the stick by grasping it from outside the bag. For best results, pull from the middle of the fork-like end, using your thumb and index finger to move the sponge back and forth until it gently releases from the stick.



7 Use the attached tie to close the Twirl-Tie™ bag by rolling the opening of the bag and then kinking the tie to the inner side.

Package Size	Description	Item No.
	<b>Pre-moistened with 10 ml</b>	<b>In Twirl-Tie™ bag</b>
75	Neutralizing buffer (NB)	10006157
75	Dey-Engley neutralizing broth (DE)	10006265
75	Lethen broth (LB)	10006298
75	Buffered peptone water (BPW)	10006267
75	0.9 % NaCl	10006609
75	Dry	10006300

## Sponge

- In Twirl-Tie™ bag
- 3" x 1.5" (7.5 x 3.5 cm) biocide-free cellulose sponge
- Available dry or pre-moistened with 10 ml buffer
- Bags can be opened easily with attached pull tabs
- Label area for sample identification
- Irradiated for proven sterility



## Sponge in Vial

- 3" x 1.5" (7.5 x 3.5 cm) biocide-free cellulose sponge
- Pre-moistened with 10 ml buffer
- In a 120 ml plastic screw-cap vial
- Label attached for sample identification



### Procedure:



1 Open bag or vial and remove sponge with gloved hand.



2 Collect sample by vigorously rubbing sponge on targeted area (e.g. 30 x 30 cm, 1 x 1 ft.).



3 Put sponge into the sterile sample bag (or back into the vial) and label bag with sample details. Always switch gloves between sampling sites.

Package Size	Description	Item No.	
		Sponge in Twirl-Tie™ bag	Sponge in Vial
	<b>Sponge pre-moistened with 10 ml of:</b>		
100	Neutralizing buffer (NB)	10006293	
100	Dey-Engley neutralizing broth (DE)	10006264	
100 (98 in Vial)	Lethen broth (LB)	10006299	10004592*
100	Buffered peptone water (BPW)	10006266	
100	0.9 % NaCl	10006493	
250	Dry	10006301	

\*other buffers available on request depending on quantities

## Fabric Swab

- 28 x 35 cm blue biocide-free fabric swab (swab cloth)
- Pre-moistened with 10 ml buffer
- In 55 oz.; 7.5 x 12" (19 x 30 cm) Twirl-Tie™ bag
- Bags can be opened easily with attached pull-tabs
- Label area for sample identification
- Irradiated for proven sterility



### Procedure:



1 Open the bag and remove fabric swab with gloved hand.



2 Collect sample by vigorously rubbing sponge on targeted area (e.g. 30 x 30 cm, 1 x 1 ft.).



3 Put the swab into the sterile Twirl-Tie™ bag and label bag with sample details. Always switch gloves between sampling sites.



4 Close the Twirl-Tie™ bag with the wires attached.

### Fabric Swabs (according to EU Commission Regulation, 28 x 35 cm, ≥ 900 cm<sup>2</sup>) in 55 oz. (1.62 L) Twirl-Tie™ bag, sterile 7.5" x 12" (19 x 30 cm)

Package Size	Description	Item No.
200	Fabric Swab pre-moistened with MRD	10004589
200	Fabric swab pre-moistened with Neutralizing buffer (NB)	10006188
200	Fabric Swab Dry	10006534

# Miscellaneous Environmental Sampling Products

## Sterile Gloves

- In convenient tear-open plastic pouch
- Packaged as a pair of gloves per pouch
- Irradiated for proven sterility



## Sterile Sampling Bag

- Sterile 18 oz (0.5 L), 24 oz (0.7 L) and 55 oz (1.62 L) bags
- Twirl-Tie™ style sample bags
- Label area for sample identification
- Bags can be opened easily with attached pull tabs
- Sterile manufactured



## Sterile Carcass Sampling Template

- 100 square cm (10 x 10 cm) or 50 square cm (10 x 5 cm)
- For carcass or surface sampling
- Made of rigid paperboard to withstand vigorous use
- Packaged in an easy-to-open plastic bag
- Irradiated for proven sterility



Package Size	Description	Item No.
100	100 cm <sup>2</sup> template packaged in re-closable bag	10003780
100	50 cm <sup>2</sup> template packaged in re-closable bag	10003786
50 pairs	Co-polymer gloves size XL packaged in peel open pouch	10002795
100	18 oz. (0.5 L) Twirl-Tie™ bag, sterile 4.5" x 9"	10001883
100	24 oz. (0.7 L) Twirl-Tie™ bag, sterile 5.5" x 9"	10001884
500	55 oz. (1.62 L) Twirl-Tie™ bag, sterile 7.5" x 12"	10001886

Photo Cover: Gettyimages@alle12